



***Andinoacara rivulatus* (Perciformes: Cichlidae), an introduced exotic fish in the upstream of Brantas River, Indonesia**

¹Novalina Serdiati, ²Danang Yonarta, ³Fajar S. Pratama, ⁴Abdul R. Faqih, ⁵Fitri S. Valen, ⁵Muhammad B. Tamam, ⁶Yogaswara I. G. Hamzah, ^{5,7}Veryl Hasan

¹ Fisheries Department, Faculty of Animal Husbandry and Fisheries, University of Tadulako, Palu, Central Sulawesi, Indonesia; ² Study Programe of Aquaculture, Faculty of Agriculture, University of Sriwijaya, Ogan Ilir, South Sumatra, Indonesia; ³ Ministry of Marine Affairs and Fisheries Republic of Indonesia, Direktorat General of Marine and Fisheries Resources Surveillance, Tual Surveillance for Marine and Fisheries Resources Base, Tual, Maluku, Indonesia; ⁴ Water Resource Management Department, Faculty of Fisheries and Marine Science, University of Brawijaya, Malang-East Java, Indonesia; ⁵ Generasi Biologi Indonesia Foundation, Zoology Division, Gresik, East Java, Indonesia; ⁶ Post Graduate School, Department of Agribusiness, Muhammadiyah Malang University, Malang, East Java, Indonesia; ⁷ Department of Fish Health Management and Aquaculture, Faculty of Fisheries and Marine, University of Airlangga, Surabaya, East Java, Indonesia.
Corresponding author: V. Hasan, verylbp@gmail.com

Abstract. The presence of non-native fish, especially types of predators was a real threat to freshwater ecosystems in Indonesia, peculiarly for native fish which were unable to compete. The entry of invasive fish had the potential as a disease vector, uncontrolled hybridization, and damaging food webs that would have a systemic impact on the ecosystem. This study discussed the first record of the existence of Green terror, *Andinoacara rivulatus* from South America in the freshwater waters of East Java, Indonesia, precisely in the upstream of Brantas River. A description of morphological characters of a specimen is provided.

Key Words: biodiversity, freshwater fish, alien fish, gold saum, non-native.

Introduction. The loss of freshwater biodiversity could be triggered by environmental changes both naturally and due to human intervention (Colautti & McIsaac 2014; Copp et al 2005). Fish that did not survive with these conditions could experience a decline in population and even extinction (Dudgeon et al 2006). The issue that was currently developing in Indonesia was the reduction of some native fish species due to the introduction of non-native species (Webb 2007; Muchlisin 2012).

The entry of foreign fish into Indonesia was basically aiming to increase aquaculture production such as Tilapia (Ishikawa et al 2012; Hasan & Tamam 2019; Hasan et al 2019a), Common carp (Weber et al 2016) and African catfish (Vitule et al 2006), but in subsequent developments, the ornamental fish from South America, especially from Peru and Ecuador (Musilova et al 2009; Wijkmark et al 2012) enter Indonesia due to market demand. One of the ornamental fish from South America that was currently circulating in the Indonesian local market was Green terror (*Andinoacara rivulatus*).

A. rivulatus exhibits highly predatory habits and tolerance to new habitats. Because of this, *A. rivulatus* has the potential to become an invasive fish and can have negative impacts on aquatic communities through competition for food, predation and other resources (Moyle & Light 1996; Sanders et al 2014). *A. rivulatus* is generally sold in

the aquarium trade as ornamental fish and it has not been cultured openly. The aim of this paper was report the first occurrence of *A. rivulatus* in upstream of Brantas River, East Java.

Material and Method. Nineteen specimens of *A. rivulatus* were sampled on 15 October 2018 and 28 December 2019 from upstream of Brantas River, East Java, Indonesia (Figure 1), on the coordinates 8°00'44.3"S and 112°20'03.5"E (Figure 1). The fishing gear used was a small hook with bottom and as bait worms were used (Stein et al 2012). Collected specimens were fixed in 10% formalin solution (Hasan et al 2019b) and deposited at the Zoology Laboratory, Generasi Biologi Indonesia, Gresik, Indonesia (GBI0023). The diagnostic of the morphological characters of the specimens were analyzed following Wijkmark et al (2012).



Figure 1. Upstream of Brantas River, East Java, the location where *A. rivulatus* was collected (original).

Results. The specimens of *A. rivulatus* collected from the upstream of Brantas River had total length between 88.1-28.7 cm (Figure 2) and weight between 944.1-340.5 g. The sample included nine females, four males and six unsex. The specific morphological characters were the follows: compressed body, short head, medium snout, dorsal fin base slightly curved, prepelvic contour straight curved, terminal mouth, scales on gill cover, cheek and prepelvic area cycloid, rounded caudal fin. Colour pattern in fresh specimens: snout and front dorsally greyish or brownish. Sides of head with 2 narrow dark lines from orbital margin to mouth. Dark spots scattered over gill cover. Dorsal, pelvic and anal fins brownish. Pectoral fin hyaline. Caudal fin dark brown mouth.



Figure 2. Male specimen of *Andinoacara rivulatus* GBI0023 captured in December 2019 from upstream of Brantas River, East Java, Indonesia (original).

The existence of *A. rivulatus* in the upstream of Brantas River is the new record of this species from East Java (Figure 3). This fact is a warning to other streams that have many native freshwater fish in such as midstream and downstream to prevent the entry of *A. rivulatus* into river basin. Water conditions in the upstream of Brantas River, (DO 7.3-9.6, temperature 22–24°C, depth 0.4–3.2, pH 7.1-7.7) are ideal for *A. rivulatus* reproduction and survival (Riehl & Baensch 1991). The size of sampled specimens, ranging from sub-adults to adults, suggest that *A. rivulatus* are able to adapt well in the upstream of Brantas River, and can possibly establish a viable population there.



Figure 3. The red square marks the location of Brantas River upstream.

There are many native aquatic organisms in the upstream of Brantas River, mostly fish (Cyprinidae, Synbranchidae, Bagridae, Sisoridae and Clariidae), that can be affected by *A. rivulatus*, including through competition for niche and predation, so that the presence of *A. rivulatus* is a danger alert.

In Indonesia, *A. rivulatus* is included in the invasive fish category, according to the Ministry of Maritime Affairs and Fisheries Regulation Number 41/2014. This information should be immediately mediatized to the local community and ornamental fish hobbyist then immediately enforced, subject to a punishment for these violators. The way to overcome alien fish that has already entered is by catching it, making it fish meal as fish feed or community food. This habit can suppress *A. rivulatus* populations so that the population is not massive.

Conclusions. The results of the identification of the morphology of alien fish in the upstream of Brantas River are *A. rivulatus*, predatory fish from South America. The existence of *A. rivulatus* in Indonesia can disturb native fish population. Cooperation between the government and the public are needed so that *A. rivulatus* to not spread to other freshwaters.

Acknowledgements. We thank to the reviewers and editors for their insightful comments. Local fishermen and Generasi Biologi Indonesia and local angler as field

guide, and the Ministry of Finance of the Indonesian for the funding of research (no. 20160221035555).

References

- Colautti R. I., MacIsaac H. L., 2004 A neutral terminology to define invasive species. *Diversity and Distributions* 10(2):135-141.
- Copp G. H., Bianco P. G., Bogutskaya N. G., Erős T., Falka I., Ferreira M. T., Fox M. G., Freyhof J., Gozlan R. E., Grabowska J., Kováč V., Moreno-Amich R., Naseka A. M., Peňáz M., Povž M., Przybylski M., Robillard M., Russell I. C., Stakénas S., Šumer S., Vila-Gispert A., Wiesner C., 2005 To be, or not to be, a non-native freshwater fish? *Journal of Applied Ichthyology* 21(4):242-262.
- Dudgeon D., Arthington A. H., Gessner M. O., Kawabata Z., Knowler D. J., Lévêque C., Naiman R. J., Prieur-Richard A. H., Soto D., Stiassny M. L., Sullivan C. A., 2006 Freshwater biodiversity: importance, threats, status and conservation challenges. *Biological Reviews of the Cambridge Philosophical Society* 81(2):163-182.
- Hasan V., Tamam M. B., 2019 First record of the invasive Nile Tilapia, *Oreochromis niloticus* (Linnaeus, 1758) (Perciformes, Cichlidae), on Bawean Island, Indonesia. *Check List* 15(1):225-227.
- Hasan V., Mukti A. T., Putranto T. W. C., 2019a Range expansion of the invasive Nile tilapia *Oreochromis niloticus* (Perciformes: Cichlidae) in Java Sea and first record for Kangean Island, Madura, East Java, Indonesia. *Ecology, Environment and Conservation* 25 (July Suppl. Issue):S187-S189
- Hasan V., Soemarno, Widodo M. S., Wiadnya D. G. R., Mukti A. T., Irawan B., 2019b Distribution extension and first record of *Lobocheilos falcifer* (Cypriniformes, Cyprinidae) in Central Java Province, Indonesia. *Ecology, Environment and Conservation* 25 (July Suppl. Issue):S119-S122.
- Ishikawa T., Shimose T., Tachihara T., 2012 Life history of an invasive and unexploited population of Nile tilapia (*Oreochromis niloticus*) and geographical variation across its native and non-native ranges. *Environmental Biology of Fishes* 96(5):603-616.
- Moyle P. B., Light T., 1996 Biological invasions of fresh water: empirical rules and assembly theory. *Biological Conservation* 78:149-161.
- Muchlisin Z. A., 2012 First report on introduced freshwater fishes in the waters of Aceh, Indonesia. *Archives of Polish Fisheries* 20:129-135.
- Musilova Z., Schindler I., Staeck W., 2009 Description of *Andinoacara stalsbergi* sp. (Teleostei: Cichlidae: Cichlasomatini) from Pacific coastal rivers in Peru, and annotations on the phylogeny of the genus. *Vertebrate Zoology* 59(2):131-141.
- Riehl R., Baensch H. A., 1991 *Aquarien atlas*. Band. 1. Melle: Mergus, Verlag für Natur- und Heimtierkunde, Germany.
- Sanders S., Astiglione C., Hoff M., 2014 Risk assessment mapping program: RAMP. U.S. Fish and Wildlife Service.
- Stein J. A., Shultz A. D., Cooke S. J., Danylchuk A. J., Hayward K., Suski C. D., 2012 The influence of hook size, type, and location on hook retention and survival of angled bonefish (*Albula vulpes*). *Fisheries Research* 113:147-152.
- Vitule J. R. S., Umbria S. C., Aranha J. M. R., 2006 Introduction of the African catfish *Clarias gariepinus* (Burchell, 1822) into Southern Brazil. *Biological Invasion* 8: 677. <https://doi.org/10.1007/s10530-005-2535-8>.
- Weber M. J., Hennen M. J., Brown M. L., Lucchesi D. O., Sauver T. R. S., 2016 Compensatory response of invasive common carp *Cyprinus carpio* to harvest. *Fisheries Research* 179:168-178.
- Webb A. C. 2007 Status of non-native freshwater fishes in tropical northern Queensland, including establishment success, rates of spread, range and introduction pathways. *Journal & Proceedings of the Royal Society of New South Wales* 140:63-78.
- Wijkmark N., Sven O. K., Salzar R. E. B., 2012 *Andinoacara blombergi*, a new species from the río Esmeraldas basin in Ecuador and a review of *A. rivulatus* (Teleostei: Cichlidae). *Ichthyological Exploration of Freshwaters* 23(2):117-137.

*** Ministry of Maritime Affairs and Fisheries, 2014 Prohibition of importing dangerous fish types from overseas into the state of the Republic of Indonesia. Regulation No. 41, Indonesia.

Received: 05 December 2019. Accepted: 14 January 2020. Published online: 23 January 2020.

Authors:

Noalina Serdiati, University of Tadulako, Faculty of Animal Husbandry and Fisheries, Fisheries Department, Indonesia, Central Sulawesi, 94118 Palu, Kampus Bumi Tadulako, e-mail: novalinaserdiati@untad.ac.id

Danang Yonarta, University of Sriwijaya, Faculty of Agriculture, Study Programe of Aquaculture, Indonesia, South Sumatra, Ogan Ilir, Palembang-Prabumulih Street, e-mail: danangyonarta@unsri.ac.id

Fajar Surya Pratama, Ministry of Marine Affairs and Fisheries Republic of Indonesia, Direktorat General of Marine and Fisheries Resources Surveillance, Tual Surveillance for Marine and Fisheries Resources Base, Indonesia, Maluku, 97651 Tual, e-mail: fajarsuryapratama@gmail.com

Abdul Rahem Faqih, University of Brawijaya, Faculty of Fisheries and Marine Science, Water Resource Management Department, Indonesia, East Java, 65145 Malang, Jl. Veteran, e-mail: faqihmadura_67@hotmail.com

Fitri Sil Valen, Generasi Biologi Indonesia Foundation, Zoology Division, Indonesia, East Java, 61171 Gresik, e-mail: valen@genbinesia.or.id

Muhammad Badrut Tamam, Generasi Biologi Indonesia Foundation, Zoology Division, Indonesia, East Java, 61171 Gresik, e-mail: tamam@genbinesia.or.id

Yogaswara Indra Ghardika Hamzah, Muhammadiyah Malang University, Department of Agribusiness, Post Graduate School, Indonesia, East Java, 65144 Malang, e-mail: yogaikhlash66@gmail.com

Veryl Hasan, University of Airlangga, Faculty of Fisheries and Marine, Department of Fish Health Management and Aquaculture, Indonesia, East Java, Surabaya 60115, Campus C, e-mail: verylbp@gmail.com

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Serdiati N., Yonarta D., Pratama F. S., Faqih A. R., Valen F. S., Tamam M. B., Hamzah Y. I. G., Hasan V., 2020 *Andinoacara rivulatus* (Perciformes: Cichlidae), an introduced exotic fish in the upstream of Brantas River, Indonesia. *AAFL Bioflux* 13(1):137-141.