

Effects of formulated diets supplemented with vitamin E on the egg quality and ovi somatic index of female *Portunus pelagicus* broodstock

¹Efrizal, ¹Indra J. Zakaria, ²Rusnam

¹ Department of Biology, Faculty of Mathematics and Natural Sciences, Andalas University, Padang, West Sumatera, Indonesia; ² Agricultural Engineering, Faculty of Agricultural Technology, Andalas University, Padang, West Sumatera Indonesia.
Corresponding author: Efrizal, e-mail: efrizal@sci.unand.ac.id

Abstract. The study of the blue swimming crab (*Portunus pelagicus*) broodstock nutrition is significant for achieving the reproductive success, which implicitly increases the larval quantity and quality in the mass production of cultured species. Therefore, the objective of this work was to determine the effects of the dosage of vitamin E in the diets on the egg quality and ovi somatic index of female *P. pelagicus* broodstock. Four dosage levels of vitamin E were tested: 0 IU kg⁻¹ (control), 300 IU kg⁻¹, 600 IU kg⁻¹, and 900 IU kg⁻¹, in a formulated diet (44.38% crude protein). *P. pelagicus* was fed once a day between 17.00 and 18.00 hours, at the ration of 3% of the body weight, for 40 days. The results showed that the fertilization rate, the egg hatching rate and the ovi somatic index increased with an increasing vitamin E dosage in the diet, from control (0 IU kg⁻¹) to 300 IU kg⁻¹, then it decreased when the dosage was increased up to 900 IU kg⁻¹ formulated diet. A quadratic relationship existed between the dose of vitamin E (IU kg⁻¹) in the formulated diet and the egg quality incubation period (days) and ovi somatic index of the female *P. pelagicus* broodstock. The modification of the formulated diet by supplementation with different doses of vitamin E can reduce the incubation period of female *P. pelagicus* broodstock and increase the egg quality and ovi somatic index value of the female parent.

Key Words: fertilization rate, fecundity, hatching rate, incubation period, nutrition.

Introduction. *Portunus pelagicus* represents a new commodity in aquaculture, with excellent development prospects. However, the breeding technology of the *P. pelagicus* cultivation is still in progress. In order to take advantage of the opportunity of an increasing market value and demand, most crab cultivators bring the seed from the wild to the culture ponds (Efrizal et al 2019a). The blue swimming crab, particularly the genus *Portunus*, is a commercially valuable species cultivated in several Asian countries, such as India, Indonesia, Philippines, Malaysia, Taiwan, China, Vietnam, and Sri Lanka. The worldwide export of the pasteurized *P. pelagicus* has a positive trend due to the main markets: the United States, Japan and Singapore, delivering a multi-million dollars annual income to Indonesia (Kangas 2000; Fahmi et al 2015; Efrizal 2017).

Currently, in Indonesia and Thailand, the culture methods for breeding, nursing and rearing the *P. pelagicus* have attained significantly higher productivity and survival rates. Although the necessary information for the development of the mass production of crab seeds is available (Soundarapandian et al 2007; Oniam et al 2012; Efrizal & Rusnam 2017; Efrizal et al 2019b) a systematic research on the optimal nutrition for the crab production systems is essential for the future breeding or culturing programs. In this study, the effects of formulated diets were examined, by evaluating the impact of different doses of vitamin E on the egg quality and ovi somatic index (OSI) of female *P. pelagicus* broodstock.

Material and Method

Time and site. The experiments were located at the Fish Seed Center Beaches (BBIP) Teluk Buo, Bungus Hatchery of Padang City. The sample analysis was carried out using

the facilities of the animal physiology laboratory, Department of Biology, Andalas University, Padang, West Sumatera, Indonesia.

Crab collection and husbandry. *P. pelagicus* in the study were obtained on January 2018. Twenty parental crabs were at the ovarian maturation stage II, with a bodyweight ranging between 170.81 and 203.53 g. The *P. pelagicus* sample specimens were captured from the wild and kept at the experimental site location. They were distributed randomly into 200 x 100 x100 cm (four concrete tanks), holding five units per tank, in a plastic box (45.5 x 32.5 x 16.5 cm), at the density of one crab per box. Tanks were equipped with hides (PVC pipes with a diameter of 13 cm and a length of 40 cm), providing a substrate of around 15 cm thick layer and an appropriate aeration (Efrizal 2015; Efrizal et al 2015; Efrizal et al 2019c). The crabs were held a pH reading of 7.26 to 8.00, an oxygen reading of 6.15 to 7.45 mg L⁻¹, a salinity reading of 29 to 32 ng L⁻¹, a temperature reading of 26-28°C and a water depth reading of 25-30 cm.

Supplementation. *P. pelagicus* was fed once a day, between 17.00 and 18.00 hours, at the ration of 3% of the body weight, for 40 days. Uneaten food was removed every morning. The experiment used a completely randomized design (CRD) consisting in 4 treatments with dietary vitamin E and 5 replications: formulated diet Fdiet 1, with 0 IU kg⁻¹; formulated diet Fdiet 2, with 300 IU kg⁻¹; formulated diet Fdiet 3, with 600 IU kg⁻¹; and formulated diet Fdiet 4, with 900 IU kg⁻¹. The diet is a moderated formulation for *Scylla serrata* broodstock (Millamena & Qunitio 2000) and *P. pelagicus* broodstock (Efrizal & Rusnam 2017; Efrizal et al 2019a; Efrizal et al 2019c). *P. pelagicus* broodstock specimens were initially fed with fresh feed, before gradually switching to the artificial experimental diet. The first ten days they were fed to satiation.

Variables assessed. Measured parameters of the egg quality (incubation period, egg diameter, fertilization rate, hatching rate, fecundity) and ovi somatic index were determined according to Efrizal et al (2006). The egg incubation period was the number of days from spawning to hatching (Efrizal et al 2006). The fertilization rate (Cerda et al 1994; Millamena & Qunitio 2000) was calculated as follows:

$$FR = FE / (FE + UFE) \times 100\%$$

Where:

FR - the fertilization rate (%);

FE - the fertilized eggs number;

UFE - the unfertilized eggs number.

The hatching rate (Millamena & Qunitio 2000; Adebayo & Popoola 2008) and fecundity (Cerda et al 1994; Oniam & Taparhudee 2010) were calculated as follows:

$$HR = HZ / (HZ + UFE) \times 100\%$$

$$F = HZ + UFE$$

Where:

HR - the hatching rate (%);

F - fecundity;

HZ - the hatched zoea number;

UFE - the unfertilized eggs number.

The ovi somatic index was calculated as follows (modified from Aryani & Suharman 2015):

$$OSI = OEW / WWC \times 100\%,$$

Where:

OSI - the ovi somatic index (%);

OEW - the ovulated egg weight (g);

WWC - the wet weight of the crab (g).

The ovulated egg weight was obtained from differences between the weight of the berried female and the weight of the non-berried female (post-spawning). Weights were measured with a precision of 0.01 g on the electronic balances (BL3200H-SHIMADZU).

The water quality parameters were determined according to Rice et al (2012) and Efrizal et al (2019c).

Statistical analysis. The data for the egg quality (incubation period, egg diameter, fertilization rate, hatching rate, and fecundity) and ovi somatic index were subjected to the analysis of variance (ANOVA), followed by Duncan's Multiple Range test to compare the mean differences among treatments (Steel & Torrie 1990). Arcsine transformation was done in the analysis of the data in percentages. The relationships among the dose of vitamin E (IU kg⁻¹) in a formulated diet and the egg quality and ovi somatic index were analyzed using the statistical software, SPSS version 19.0, to identify significant correlations between them.

Results and Discussion

Egg quality

Incubation period. The data (Table 1) show that the incubation period was not significantly ($P > 0.05$) affected by the formulated diets of vitamin E supplementation at different doses. The doses of vitamin E of 300 IU kg⁻¹ in the formulated diet resulted in the shortest incubation period (6.80±0.42 days). The longest incubation period was achieved at Fdiet 1 (vitamin E 0 IU kg⁻¹), Fdiet 3 (vitamin E 600 IU kg⁻¹) and Fdiet 4 (vitamin E 900 IU kg⁻¹) with 7.60±0.27, 7.20±0.27 and 7.20±0.27 days, respectively. The relationship between the doses of vitamin E (DVE) in the formulated diet and the incubation period (IP) is shown in Figure 1. The regression equation is $IP = 3 \times 10^{-6} DVE^2 - 0.0026 DVE + 7.53$ ($R^2 = 0.1626$; $P > 0.05$).

Table 1
The incubation period (days) of female *Portunus pelagicus* broodstock at a different dosage of supplemented vitamin E in the formulated diet

Treatment (n=5)	The incubation period (days)
Fdiet 1	7.60±0.27 ^a
Fdiet 2	6.80±0.42 ^a
Fdiet 3	7.20±0.22 ^a
Fdiet 4	7.40±0.27 ^a

Means (±SE) within a given column with different superscripts are significantly different ($P < 0.05$); n-replication; Fdiet 1-vitamin E 0 IU kg⁻¹ formulated diet (control); Fdiet 2-vitamin E 300 IU kg⁻¹ formulated diet; Fdiet 3-vitamin E 600 IU kg⁻¹ formulated diet; Fdiet 4-vitamin E 900 IU kg⁻¹ formulated diet.




Figure 1. The relationship between the dose of vitamin E (IU kg⁻¹) in a formulated diet and the incubation period (days) of female *Portunus pelagicus* broodstock.

Egg diameter. The data on the egg diameter are summarized in Table 2 and Figure 2. Viable eggs of berried female *P. pelagicus* were spherical, yellowish-orange, and

somewhat translucent, and ranged from 0.334-0.357 mm in diameter. The greatest egg diameter (0.357 mm) of the berried female was observed in the treatment with 300 IU kg⁻¹ vitamin E in the formulated diet, and the lowest egg diameter was found in the treatment of 0 IU kg⁻¹ vitamin E in the formulated diet (Table 2). The ANOVA showed that the treatment significantly (P<0.05) affected the egg diameter of the berried female blue swimming crab, and then, Duncan's test showed that the Fdiet 2 effect was significantly different (P<0.05) compared to that of Fdiet 1 on the egg diameter. The relationship between the doses of vitamin E in the formulated diet and the egg diameter (ED) is shown in Figure 2. The regression equation was found to be quadratic and was calculated as follows: $ED = -8 \times 10^{-8} DVE^2 + 9 \times 10^{-5} DVE + 0.3347$. The $R^2 = 0.5772$ (P>0.05) indicated an increase of the egg diameter of approximately 42% with the addition of vitamin E to the formulated diet (IU kg⁻¹).

Table 2
Mean egg diameter (mm) of female *Portunus pelagicus* broodstock, at a different dosage of supplemented vitamin E in formulated diet

Replication (n=5)	Treatment			
	Fdiet 1	Fdiet 2	Fdiet 3	Fdiet 4
1	0.333±0.031	0.360±0.026	0.354±0.020	0.342±0.019
2	0.327±0.024	0.350±0.029	0.350±0.021	0.355±0.020
3	0.338±0.026	0.353±0.027	0.355±0.023	0.350±0.025
4	0.335±0.029	0.369±0.030	0.346±0.028	0.351±0.026
5	0.335±0.029	0.355±0.034	0.377±0.023	0.365±0.020
ED (mm)	0.334±0.002 ^a	0.357±0.004 ^b	0.356±0.006 ^b	0.353±0.004 ^b

Means (±SE) within a given column with different superscripts are significantly different (P<0.05); n-replication; ED-egg diameter; Fdiet 1-vitamin E 0 IU kg⁻¹ formulated diet (control); Fdiet 2-vitamin E 300 IU kg⁻¹ formulated diet; Fdiet 3-vitamin E 600 IU kg⁻¹ formulated diet; Fdiet 4-vitamin E 900 IU kg⁻¹ formulated diet.




Figure 2. The relationship between the dose of vitamin E (IU kg⁻¹) in a formulated diet and the egg diameter (mm) of female *Portunus pelagicus* broodstock.

Fertilization rate. The success rate for egg fertilization with the different diet treatments was generally high and ranged from 80.53 to 96.42% (Table 3). The fertilization rate (%) decreased with the increasing dose of vitamin E in the formulated diet (P<0.05). The highest fertilization rate (96.42±0.98%) was achieved at the dose of vitamin E of 300 IU kg⁻¹ in the formulated diet, whereas the lowest (P<0.05) fertilization rate occurred at the dose of vitamin E of 0 IU kg⁻¹ in the formulated diet with 80.53±0.73% (Table 3). The relationship between the dose of vitamin E in the formulated diet and the egg fertilization rate (FR) was quadratic, with an equation of $FR = -5 \times 10^{-5} DVE^2 + 0.0505 DVE + 81.81$ (Figure 3). This relationship was found to be highly significant ($R^2 = 0.6584$; P<0.05).

Table 3

Mean egg fertilization rate (%) of female *Portunus pelagicus* broodstock, at a different dosage of supplemented vitamin E in formulated diet

Treatment (n=5)	Number of eggs sampled		Fertilization rate (%)
	Fertilized	Total	
Fdiet 1	534.40±2.02	688.60±6.34	80.53±0.73 ^a
Fdiet 2	656.40±11.82	680.00±10.23	96.42±0.98 ^b
Fdiet 3	619.20±12.58	682.20±10.09	90.75±0.94 ^c
Fdiet 4	607.20±15.68	681.00±6.79	89.14±1.76 ^c

Means (±SE) within a given column with different superscripts are significantly different (P<0.05); n-replication; Fdiet 1-vitamin E 0 IU kg⁻¹ formulated diet (control); Fdiet 2-vitamin E 300 IU kg⁻¹ formulated diet; Fdiet 3-vitamin E 600 IU kg⁻¹ formulated diet; Fdiet 4-vitamin E 900 IU kg⁻¹ formulated diet.




Figure 3. The relationship between the dose of vitamin E (IU kg⁻¹) in a formulated diet and the fertilization rate (%) of female *Portunus pelagicus* broodstock.

Hatching rate. The formulated diet with vitamin E supplementation (0, 300, 600, and 900 IU kg⁻¹ formulated diet) significantly influenced (P<0.05) the hatching rate (Table 4).

Table 4

Mean zoeal hatching rate (%) of female *Portunus pelagicus* broodstock at a different dosages of supplemented vitamin E in formulated diet

Treatment (n=5)	Number of eggs			Hatching rate (%)
	Hatching	Unfertilized	Total	
Fdiet 1	464,560.42± 118,909.36	128,527.20± 33,654.05	593,087.62± 152,108.72	78.29±0.83 ^a
Fdiet 2	668,294.51± 136,861.03	86,448.06± 17,363.87	754,742.57± 151,107.22	88.46±1.08 ^b
Fdiet 3	577,812.58± 119,869.02	281,853.10± 111,083.11	688,895.68± 146,080.61	84.05±0.92 ^c
Fdiet 4	546,864.36± 118,617.25	116,287.65± 26,609.22	663,152.01± 144,602.41	82.48±0.94 ^c

Means (±SE) within a given column with different superscripts are significantly different (P<0.05); n-replication; Fdiet 1-vitamin E 0 IU kg⁻¹ formulated diet (control); Fdiet 2-vitamin E 300 IU kg⁻¹ formulated diet; Fdiet 3-vitamin E 600 IU kg⁻¹ formulated diet; Fdiet 4-vitamin E 900 IU kg⁻¹ formulated diet.

Similar to the fertilization rate, the highest zoeal hatching rate of *P. pelagicus* eggs was achieved in the groups of crab fed on dietary treatments at 300 IU kg⁻¹ vitamin E in formulated diet (88.46±1.08%), whereas the lowest hatching rate of eggs was observed at 0 IU kg⁻¹ vitamin E in formulated diet (78.29±0.83%), compared to Fdiet 3 (300 IU kg⁻¹ vitamin E in formulated diet; 84.05±0.92%) and Fdiet 4 (300 IU kg⁻¹ vitamin E in formulated diet; 82.48±0.94%). Duncan's test demonstrated that the zoeal hatching rate

reared at the 0 IU vitamin E kg⁻¹ formulated diet was significantly lower than those reared with the other formulated diets.

The relationship between the zoal hatching rate and dose of vitamin E in the formulated diet was found to be quadratic ($HR = -3 \times 10^{-5} DVE^2 + 0.0321 DVE + 79.157$, $R^2 = 0.5864$, $P < 0.05$) (Figure 4).




Figure 4. The relationship between the dose of vitamin E (IU kg⁻¹) in a formulated diet and the hatching rate (%) of female *Portunus pelagicus* broodstock.

Fecundity. The estimated mean number of eggs per batches produced by female crabs with the different formulated diets after hatching under laboratory conditions ranged from 593,087.62 to 754,742.57 eggs batch⁻¹ (Table 5). The lowest fecundity (593,087.62±152,108.72 eggs batch⁻¹) was found in a berried female *P. pelagicus* that received the formulated diet supplementation of vitamin E 0 IU kg⁻¹ (Fdiet 1), whereas the highest fecundity (754,742.57±151,107.22 eggs batch⁻¹) was found in a berried female *P. pelagicus* receiving the formulated diet supplementation of vitamin E 300 IU kg⁻¹ (Fdiet 2). On the other hand, females fed with Fdiet 3 (600 IU kg⁻¹ formulated diet) and Fdiet 4 (900 IU kg⁻¹ formulated diet) showed a decreased fecundity, 688,895.68±146,080.61 and 663,152.01±144,602.41 eggs batch⁻¹, compared to those fed with Fdiet 2 (300 IU kg⁻¹ formulated diet), but the results of the variance tests (ANOVA) for this difference did not reveal a significant difference ($P > 0.05$). The relationship between the fecundity and dose of vitamin E in the formulated diet was quadratic with an equation of $F = -0.5206 DVE^2 + 516.61 DVE + 606468$ ($R^2 = 0.0332$, $P > 0.05$) (Figure 5).

Table 5
Mean fecundity (egg batch⁻¹) of female *Portunus pelagicus* broodstock, at a different dosage of supplemented vitamin E in formulated diet

Replication (n=5)	Treatment			
	Fdiet 1	Fdiet 2	Fdiet 3	Fdiet 4
1	358,389.72	606,186.79	499,408.48	504,048.09
2	1,068,355.99	500,062.22	917,012.05	1,178,255.90
3	384,044.25	559,902.79	568,548.58	569,488.84
4	727,380.36	878,290.59	1,073,024.15	551,069.52
5	427,267.77	1,229,270.45	386,485.15	512,897.71
F	593,087.62± 152,108.72 ^a	754,742.57± 151,107.22 ^a	688,895.68± 146,080.61 ^a	663,152.01± 144,602.41 ^a

Means (±SE) within a given column with different superscripts are significantly different ($P < 0.05$); n-replication; F-fecundity; Fdiet 1-vitamin E 0 IU kg⁻¹ formulated diet (control); Fdiet 2-vitamin E 300 IU kg⁻¹ formulated diet; Fdiet 3-vitamin E 600 IU kg⁻¹ formulated diet; Fdiet 4-vitamin E 900 IU kg⁻¹ formulated diet.




Figure 5. The relationship between the dose of vitamin E (IU kg⁻¹) in a formulated diet and the fecundity (egg batch⁻¹) of female *Portunus pelagicus* broodstock.

Ovi somatic index. The final percentage for the ovi somatic index (OSI) differed significantly ($P < 0.05$) between treatments (Table 6). This index ranged from 7.67-22.94% in all the experiments. The highest OSI value of broodstock females in this study was obtained from females receiving the Fdiet 2 and was 22.94% (Table 6 and Figure 5), followed by Fdiet 3 (17.09%), Fdiet 4 (14.77%) and Fdiet 1 (7.67%). When further tested with the Duncan's test, a significant difference ($P < 0.05$) was also observed among Fdiet 1, Fdiet 2 and Fdiet 3, no significant difference ($P > 0.05$) was observed between Fdiet 1 and Fdiet 4 (Table 6). The R^2 , which was of 0.3884 ($OSI = -5 \times 10^{-5} DVE^2 + 0.0491 DVE + 8.9086$, $P > 0.05$) (Figure 6), suggested that approximately 62% of *P. pelagicus* showed an effect in the OSI with the addition of vitamin E in the formulated diet (IU kg⁻¹).

Table 6
Mean ovi somatic index (%) of female *Portunus pelagicus* broodstock, at a different dosage of supplemented vitamin E in the formulated diet

Treatment (n=5)	Body weight before berried (g)	Body weight after berried (g)	Egg batch weight (g)	Ovi somatic index (%)
Fdiet 1	170.81±30.02	158.73±30.19	12.08±1.42	7.67±1.27 ^a
Fdiet 2	203.53±27.87	158.77±26.93	44.76±3.19	22.94±2.75 ^b
Fdiet 3	191.15±28.36	158.75±25.62	32.40±6.50	17.09±3.65 ^b
Fdiet 4	186.68±26.98	159.32±24.37	27.35±6.43	14.77±3.45 ^{ab}

Means (±SE) within a given column with different superscripts are significantly different ($P < 0.05$); n-replication; Fdiet 1-vitamin E 0 IU kg⁻¹ formulated diet (control); Fdiet 2-vitamin E 300 IU kg⁻¹ formulated diet; Fdiet 3-vitamin E 600 IU kg⁻¹ formulated diet; Fdiet 4-vitamin E 900 IU kg⁻¹ formulated diet.




Figure 6. The relationship between the dose of vitamin E (IU kg⁻¹) in a formulated diet and the ovi somatic index (%) of female *Portunus pelagicus* broodstock.

Discussion. The egg incubation period of *P. pelagicus* decreased with an increasing dose of vitamin E in the formulated diet. However, high doses of vitamin E had an inverse effect and inhibited the egg incubation period. This phenomenon was observed in the treatments with the 600-900 IU kg⁻¹ vitamin E in the formulated diet, with the egg incubation period of berried female of the *P. pelagicus* being longer compared to those females receiving the treatment of 300 IU/kg vitamin E in the formulated diet. The difference in the achievement time of the incubation period is thought to be due to differences in the quality of the diet given, which affects the quality of the eggs produced. Vitamin E plays an essential role in gonad development, namely for the process of fertilization and influences fecundity (Izquierdo et al 2001). Vitamin E can be added to feed to accelerate the phase of follicular formation (Verakunpiriya et al 1997). Vitamin E is transported from the peripheral tissues during vitellogenesis even though the plasma content of the vitellogenin is not affected, and it is suspected that lipoproteins might be involved in the transport of vitamin E during the vitellogenesis (Izquierdo et al 2001).

Compared to the results of the studies related to the time of incubation of crustaceans under the influence of other environmental factors such as temperature, the control of the environmental factors in the diet, as in the current study, is more capable of accelerating a decrease in the *P. pelagicus* egg incubation period. Such control was shown by the results reported by Hamasaki et al (2002) who found that controlling the temperature in the range from 20.3-30°C is only able to reduce the incubation period for 10-30 days in *S. serrata*. For the same species, *P. pelagicus*, the incubation period was found to range from 8.33-6.67 days at 28-34°C (Efrizal et al 2006).

The differences in the egg diameters observed in this study and analyzed quantitatively are presumably attributable to the biological response to the quality of feed given to the target cells. In this case, the feed stimulated the process of biosynthesis of vitellogenesis and stimulated the germinal epithelium in the secretion of maturation-inducing steroids (MIS) that stimulate the hydration of eggs. For example, a low protein, high-calorie diet caused a reduction in red seabream (*Pagellus bogaraveo*) reproductive performance (Watanabe et al 1984). In another sparid, the gilthead seabream (*Sparus aurata*), a well-balanced broodstock diet in essential amino acids, improved the vitellogenin synthesis (Tandler et al 1995). Moreover, a reduction in the dietary protein levels from 51 to 34%, together with an increase in dietary carbohydrate levels from 10 to 32%, reportedly reduced egg viability in seabass (Cerda et al 1994). These diets have been shown to cause alterations in GnRH release in seabass broodstock during spawning (Kah et al 1994) and in the hormonal plasma levels of the gonadotropin GtH II. The latter is known to play an essential role in oocyte maturation and ovulation (Navas et al 1996).

As shown in Table 3, a significant difference ($P < 0.05$) can be seen between Fdiet 1 and Fdiets 2, 3, and 4. The difference in the percentage value of the degree of fertilization of the eggs between treatments was related to the incubation period, and to the difference in egg diameter obtained, as described above. This finding is in line with the results of the study by Efrizal et al (2012), who reported that the degree of egg fertilization of the crab female parent decreased with the length of the egg incubation time. Furthermore, one of the nutrients known to be necessary for the fertilization is vitamin E (Izquierdo & Fernandez-Palacios 1997; Izquierdo et al 2001). Vitamin E has been shown to play an essential role in reproduction (Watanabe et al 1991). Its role as an inter- and intracellular antioxidant, maintaining homeostasis of labile metabolites in the cell and tissue plasma, is well known (Izquierdo et al 2001). The antioxidant function of vitamin E can provide an essential protective role for the sperm cells during spermatogenesis and until fertilization by reducing the risk of lipid peroxidation, which is detrimental for sperm motility (Izquierdo et al 2001).

The hatching rate with the Fdiet 1 treatment when compared with the provision of Fdiets 2, 3, and 4, was related to the low percentage of fertilization, as presented in Table 8 and Figure 4. Efrizal et al (2012) reported that the hatching rate of the crab eggs tended to be in line with the percentage of egg fertilization. Besides, the hatching rate was also related to the incubation period, and the diameter of the egg produced. Davis (1981) stated that the diameters of decapod crustacean eggs increased during the incubation period, which was due to a slow enlargement of the perfect osmotic inner egg

membrane or to an enlargement of the embryo itself. Furthermore, an increase in size resulted in the absorption of water by increasing the solution concentration inside the egg until hatching time.

Besides, the mean hatching rates cited in this study may be relatively high (78.29-88.46%), these rates are probably derived from the vitamin E in the formulated diet. Vitamin E is a potent scavenger of active oxygen species and has been shown to have a protective role against the action of free radicals. Free radicals can deteriorate egg membranes and membrane integrity. Although the harmful effects of vitamin E deficiency on the reproductive performance of higher vertebrates has been demonstrated since the early 1920s, dietary vitamin E has only been shown to be an essential nutrient for fish reproduction in 1990. Its deficiency results in immature gonads in carp and *Plecoglossus altivelis*, and in reduced hatching and fry survival rates in *P. altivelis* (Watanabe 1990).

The difference in the fecundity of female crabs in quantity is thought to be the result of giving a diet supplemented with vitamin E at different doses. The difference in the dose of vitamin E in artificial feed is thought to cause changes in the vitellogenesis process in the gonad, especially in producing a balance of essential fatty acids. Furuita et al (2000) states that one of the factors that influence fecundity is the quality of feed given to test animals. Millamena & Quinitio (2000) also stated that the ratio between the levels and types of fatty acids omega-3 and omega-6 affected the fecundity of the mud crab *S. serrata*.

A difference in the OSI values was observed between dietary treatments given to the female crab parent that are allegedly related to the dose of vitamin E, which causes a balance of nutrient content in the vitellogenesis process in the gonads of the test animals. With Fdiet 2, the essential fatty acids omega-3 and omega-6 were in a more balanced composition compared to the Fdiet 3, Fdiet 4 and Fdiet 1 during the vitellogenesis process. Rusdi & Ahmad (1993) reported that the administration of a combination of fresh feed, namely, lemuru and clam meat, was the best way to increase the maturation of the gonads (100%) and to spawn (91.6%) in mangrove crabs *S. serrata*. Furthermore, lemuru and clam meat, are foods naturally rich in essential fatty acids omega-3 and omega-6, which are necessary to the process of gonadal maturation. Mokoginta (1992) explained that the levels and types of fatty acids omega-3 and omega-6 contained in the ration affected the development of the parent and the hatchability of catfish eggs. Watanabe et al (1991) reported that the lack of essential fatty acids affected the fish spawning activities.

Conclusions. The following conclusions can be drawn from the results of this experiment: (1) the manipulation of a formulated diet supplemented with different doses of vitamin E can reduce the incubation period of the female crab parent and increase the egg quality (egg diameter, fertilization rate, hatching rate, and fecundity) and ovi somatic index of the female crab parent; (2) the feeding supplemented with different vitamin E doses had a significant effect ($P < 0.05$) on the egg fertilization rate, the egg hatching rate, and the ovi somatic index (OSI), with highest values of 96.42%, 88.46%, and 22.94, respectively, with Fdiet 2; (3) the feeding supplemented with different doses of vitamin E had no significant effect ($P > 0.05$) on the incubation period (6.80-7.60 days), egg diameter (0.334-0.357 mm) and egg fecundity (593,087.62-754,742.57 egg crab⁻¹); (4) a quadratic relationship existed between the dose of vitamin E (IU kg⁻¹) in the formulated diet and the egg quality and ovi somatic index of female *P. pelagicus* broodstock.

Acknowledgements. This study was funded by the Directorate of Research and Community Service, Ministry of Research Technology and the Higher Education Republic of Indonesia (No. T/26/UN.16.17/PT.01.03/PT-PP/2019) through the National Strategic Research Institution and by the Andalas University, Research and Community Service Institution (No. T/49/UN.16.17/PP OK-KRP2GB/LPPM/2019).

References

- Aryani N., Suharman I., 2015 Effect of dietary protein level on the reproductive performance of female of green catfish (*Hemibagrus nemurus* Bagridae). *Journal of Aquaculture Research & Development* 6(11):1-5.
- Cerdá J., Carrillo M., Zanuy S., Ramos J., de la Higuera M., 1994 Influence of nutritional composition of diet on sea bass, *Dicentrarchus labrax* L., reproductive performance and egg and larval quality. *Aquaculture* 128:345-361.
- Davis C., 1981 Mechanisms of hatching in aquatic invertebrate eggs. *Oceanography and marine biology: An annual review*, Barnes 19:95-123.
- Efrizal, Arshad A., Kamarudin M. S., Saad C. R., 2006 Effects of temperature on the incubation period and reproductive performance of berried female blue swimming crab, *Portunus pelagicus* (Linnaeus, 1758) under cultured conditions. *Research Journal of Fisheries and Hydrobiology* 1(1):23-27.
- Efrizal, Zakaria I. J., Bulanin U., 2012 [Effect of combination and level of natural diets on survival rate and larval development of the blue swimming crab *Portunus pelagicus* (Linnaeus, 1758) under laboratory conditions]. In: [The role of mathematics and natural sciences in the development of human resources and natural resources]. Sipahutar H., Restuati M., Nasution Y., Silitonga M., Gultom E. S. (eds), pp. 51-58, Proceedings of the National Seminar on Seminary in the West Region State Higher Education Cooperation Agency (BKS-PTN) for Mathematics and Natural Sciences (MIPA), Faculty of Mathematics and Natural Sciences, University of North Sumatra, Medan. [In Indonesian].
- Efrizal, 2015 Effects of salinity on survival rate and larval development of blue swimming crab *Portunus pelagicus* (Linnaeus, 1758) under laboratory conditions. *Australian Journal of Basic and Applied Sciences* 9(7):557-563.
- Efrizal, Arshad A., Kamarudin M. S., Saad C. R., Amin S. M., 2015 Some aspects of reproductive biology of blue swimming crab, *Portunus pelagicus* (Linnaeus, 1758) under laboratory conditions. *Journal of Fisheries and Aquatic Science* 10(2):77-91.
- Efrizal, 2017 Effects of stocking density on survival rate and larval development of blue swimming crab, *Portunus pelagicus* (Linnaeus, 1758) under laboratory conditions. *AAFL Bioflux* 10(2):217-226.
- Efrizal, Rusnam, 2017 The Influence of dietary vitamin E on the gonad development of blue swimming crab, *Portunus pelagicus* (Linnaeus, 1758). *Australian Journal of Basic and Applied Sciences* 11(1):7-15.
- Efrizal E., Rusnam R., Suryati S., Yolanda N., Syaiful F. L., Mardiah A., 2019a Evaluation of formulated diets enriched by Amaranth extracts for the broodstock females, *Portunus pelagicus* (Linnaeus, 1758). *Pakistan Journal of Biological Sciences* 22 (6):283-290.
- Efrizal E., Syam Z., Rusnam R., Suryati S., 2019b Growth performance and survival rate of *Portunus pelagicus* (Linnaeus, 1758) broodstock females fed varying doses of amaranth extracts. *F1000 Research* 8:1466.
- Efrizal E., Zakaria I. J., Rusnam R., Suryati S., Yolanda N., 2019c Studies on biological test of formulated diets supplementation of vitamin E for the broodstock of females blue swimming crab, *Portunus pelagicus* (Linnaeus, 1758). *F1000 Research* 7:1780.
- Fahmi A. S., Maksum M., Suwondo E., 2015 USFDA Import refusal and export competitiveness of Indonesian crab in US Market. *Agriculture and Agricultural Science Procedia* 3:226-230.
- Furuita H., Tanaka H., Yamamoto T., Shirashi M., Takeuchi T., 2000 Effects of n-3 HUFA levels in broodstock diet on the reproductive performance and egg and larval quality of the Japanese flounder (*Paralichthys olivaceus*). *Aquaculture* 187:387-398.
- Hamasaki K., Suprayudi M. A., Takeuchi T., 2002 Mass mortality during metamorphosis to megalops in the seed production of mud crab *Scylla serrata* (Crustacea, Decapoda, Portunidae). *Fisheries Science* 68:1226-1232.
- Izquierdo M. S., Fernández-Palacios H., Tacon A. G. J., 2001 Effect of broodstock nutrition on reproductive performance of fish. *Aquaculture* 197:25-42.

- Izquierdo M. S., Fernández-Palacios H., 1997 Nutritional requirements of marine fish larvae and broodstock. *Cahiers OPTIONS Méditerranéennes* 22:243–264.
- Kah O., Zanuy S., Pradelles P., Cerdá J., Carrillo M., 1994 An enzyme immunoassay for salmon gonadotropin-releasing hormone and its application to the study of the effects of diet on brain and pituitary GnRH in the sea bass, *Dicentrarchus labrax*. *General and Comparative Endocrinology* 95:464–474.
- Kangas M. I., 2000 Synopsis of the biology and exploitation of the blue swimmer crab, *Portunus pelagicus* Linnaeus, in Western Australia. Fisheries Research Report No. 121, Department of Fisheries, Western Australia, 22 p.
- Millamena O. M., Qunitio E., 2000 The effects of diets on reproductive performance of eyestalk ablated and intact mud crab *Scylla serrata*. *Aquaculture* 181:81-90.
- Mokoginta I., 1992 Essential fatty acids requirements of catfish (*Clarias batrachus* Linn) for broodstock development. Dissertation, Pasca Sarjana IPB-Bogor, 80 p.
- Navas J. M., Trush M., Ramos J., Bruce M., Carrillo M., Zanuy S., Bromage N., 1996 The effect of seasonal alteration in the lipid composition of broodstock diets on egg quality in the European sea bass. *Journal of Fish Biology* 51:760-773.
- Oniam V., Chuchit L., Arkronrat W., 2012 Reproductive performance and larval quality of blue swimming crab (*Portunus pelagicus*) broodstock, fed with different feeds. *Songklanakarin Journal of Science and Technology* 34(4):381-386.
- Oniam V., Taparhudee W., 2010 Physical relationship on fecundity and hatching rate of blue swimming crab (*Portunus pelagicus* Linnaeus, 1758) from broodstocks, earthen ponds. The proceeding of 48th Kasetsart University Annual Conference, Bangkok, Thailand, pp. 99-107.
- Rice E. W., Baird R. B., Eaton A. D., Clesceri L. S., 2012 Standard methods for the examination of water and wastewater. 22nd edition. American Public Health Association (APHA), Washington D.C., USA, 1360 p.
- Rusdi I., Ahmad T., 1993 [Maturation of mud crab gonad *Scylla serrata* with various diets combinations]. Sub Balitkandita Gondol, Bali, pp. 127-132. [In Indonesian].
- Soundarapandian P., Thamizhazhagan E., Samuel N. J., 2007 Seed production of commercially important *Portunus pelagicus* (Linnaeus). *Journal of Fisheries and Aquatic Science* 2:302-309.
- Steel R. D. G., Torrie J. H., 1990 Principles and procedures of statistics. McGraw-Hill Book Company, New York, pp. 1-42.
- Tandler A., Harel M., Koven W. M., Kolkovsky S., 1995 Broodstock and larvae nutrition in gilthead seabream *Sparus aurata* new findings on its involvement in improving growth, survival and swim bladder inflation. *Israeli Journal of Aquaculture Bamidgheh* 47:95–111.
- Verakunpiriya V., Watanabe K., Mushiake K., Kawano K., Kobayashi T., Hasegawa I., Kiron V., Satoh S., Watanabe T., 1997 Effect of a krill meal supplementation in soft-pellets on spawning and quality of egg of yellowtail. *Fisheries Science* 63(3):433–439.
- Watanabe T., 1990 Effect of broodstock diets on reproduction of fish. *Actes Colloques Ifremer* 9:542–543.
- Watanabe T., Lee M., Mizutani J., Yamada T., Satoh S., Takeuchi T., Yoshida N., Kitada T., Arakawa T., 1991 Effective components in cuttlefish meal and raw krill for improvement of quality of red sea bream *Pagrus major* eggs. *Nippon Suisan Gakkaishi* 57(4):681–694.
- Watanabe T., Takeuchi T., Saito M., Nishimura K., 1984 Effect of low protein–high calorie or essential fatty acid deficiency diet on reproduction of rainbow trout. *Nippon Suisan Gakkaishi* 50(7):1207–1215.

Received: 04 February 2020. Accepted: 26 March 2020. Published online: 03 April 2020.

Authors:

Efrizal, Andalas University, Faculty of Mathematics and Natural Sciences, Department of Biology, Kampus Unand Limau Manis, Padang, 25163 West Sumatera, Indonesia, e-mail: efrizal@sci.unand.ac.id

Indra Junaidi Zakaria, Andalas University, Faculty of Mathematics and Natural Sciences, Department of Biology, Kampus Unand Limau Manis, Padang, 25163 West Sumatera, Indonesia, e-mail: indrajunaidi@sci.unand.ac.id

Rusnam, Andalas University, Faculty of Agricultural Technology, Agricultural Engineering, Kampus Unand Limau Manis, Padang, 25163 West Sumatera, Indonesia, e-mail: rusnam@ae.unand.ac.id

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Efrizal, Zakaria I. J., Rusnam, 2020 Effects of formulated diets supplemented with vitamin E on the egg quality and ovi somatic index of female *Portunus pelagicus* broodstock. AACL Bioflux 13(2):768-779.